

# Automated Lexical Analysis of Interviews with Schizophrenic Patients

Shihao Xu<sup>‡</sup>, Zixu Yang<sup>†</sup>, Debsubhra Chakraborty<sup>\*</sup>, Yasir Tahir<sup>\*</sup>, Tomasz Maszczyk<sup>‡</sup>, Victoria Yi Han Chua<sup>‡</sup>, Justin Dauwels<sup>‡</sup>, Daniel Thalmann<sup>\*</sup>, Nadia Magnenat Thalmann<sup>\*</sup>, Bhing-Leet Tan<sup>†</sup><sup>§</sup>, and Jimmy Lee Chee Keong<sup>†</sup><sup>§</sup>

**Abstract** Schizophrenia is a chronic brain disorder that contributes to poor function and quality of life. Early diagnosis and treatment of schizophrenia can effectively improve the quality of life of patients. We are aiming to design objective assessment tools of schizophrenia. In earlier work, we investigated non-verbal quantitative cues for this purpose. In this paper, we instead explore linguistic cues, extracted from interviews of schizophrenia patients and healthy control subjects, conducted by trained psychologists. Specifically, we analyzed the interviews of 47 schizophrenic patients and 24 healthy age-matched control subjects. We applied automated speech recognition and linguistic tools to capture the linguistic categories of emotional and psychological states. Based on those linguistic categories, we applied a binary classifier to distinguish patients from matched control subjects, leading to a classification accuracy of about 86% (by leave-one-out cross-validation); this result seems to suggest that schizophrenia patients tend to talk about different topics and use different words. We provided an in-depth discussion of the most salient lexical features, which may provide some insights into the linguistic alterations in patients.

## 1 Introduction

Schizophrenia is a chronic mental disorder affecting millions of people globally. Its symptoms are typically classified into three broad groups: positive (hallucinations and delusions), negative (apathy, blunting of affect, alogia), and cognitive (problems with attention, memory, and executive functions) [1]. The heterogeneous linguistic alterations associated with schizophrenia have been extensively studied since the

---

<sup>‡</sup> School of Electrical and Electronic Engineering, Nanyang Technological University, Singapore

<sup>\*</sup> Institute for Media Innovation, Nanyang Technological University, Singapore

<sup>†</sup> Institute of Mental Health, Singapore

<sup>§</sup> Health and Social Sciences, Singapore Institute of Technology

<sup>§</sup> Lee Kong Chian School of Medicine, Nanyang Technological University

last century [2, 3]. As words and expressions contain a wealth of information regarding emotions and behavior [4, 5], linguistic analysis is becoming a common tool for research in psychiatry [6, 7]. In recent years, technological advances have made it increasingly faster and more cost-effective to collect a variety of data, a data-driven approach may lead to novel discoveries and treatments in the realm of mental health [8]. In this paper, we follow such data-driven approach.

A typical lexical based tool for analyzing language is the Linguistic Inquiry and Word Count (LIWC) [9]. The latest version, LIWC 2015 [10], provides several subsets of words representing different emotional states or characteristics. Some studies have applied LIWC to essays written by schizophrenic patients. A linguistic analysis of written self-descriptive tasks completed by schizophrenic patients and controls found no differences in the use of words related to emotions (both positive and negative) between patients and control subjects when describing themselves [11]. On the other hand, analysis of standardized written assignments by psychiatric outpatients and control subjects revealed that patients used fewer words pertaining to optimism/energy, basic cognitive mechanisms, exclusion, and bodily functions [12]. In another study by Deutsch-Link et al. [13], essays written by schizophrenic patients included more external referential language and fewer mentions of the word *I* than control subjects.

LIWC has also been applied to transcripts of interviews with schizophrenic patients. A study by Minor et al. [14] consisted of analyzing structured interviews with schizophrenic patients. Interviews were recorded, manually transcribed, and then processed for lexical analysis by LIWC, where *anger* words significantly predicted greater symptoms. Similarly, Hong et al. [15] extracted lexical features to distinguish between schizophrenia patients and controls from manually transcribed speech of schizophrenia patients and control subjects. In their study, the speech of schizophrenic patients featured less usage of *I* and *adverb*, and more frequently included words from the categories of *friend* and *relative* instead, when compared to control subjects.

In all above-mentioned studies, the texts were either written by patients or manually transcribed from audio recordings. In our study, we explored the feasibility of applying linguistic analysis to automated transcriptions of interviews with patients and control subjects, conducted by psychologists. The audio recordings of 71 participants (47 patients and 24 control subjects) were automatically transcribed through speech recognition software. We then apply LIWC to the automatically transcribed text to explore linguistic differences between schizophrenic patients and controls. This research is in alignment with our previous works to develop automated, objective methods to examine behavioral deficits in schizophrenia, in which we studied non-verbal cues related to speech [16] and movement [17].

Speech recognition technology has greatly improved in recent years thanks to the breakthroughs in the domain of artificial intelligence. Many existing APIs support transcription of microphone streams and audio files directly to text files [18]. Dozens of languages are supported by these platforms, but the performance varies as they employ different language models and machine learning algorithms. Generally, it

has been observed that Deep Neural Network (DNN) based tools outperform those based on Gaussian mixture models (GMM) [19].

In our analysis, Google Cloud Speech API was utilized to convert the interview recordings to texts. By means of lexical cues, we were able to classify patients and healthy control subjects at an accuracy of 86% (by leave-one-out cross-validation). We discovered that the speech produced by schizophrenic patients featured fewer *informal* words (e.g., okay, coz, oh) and female patients speech more frequently contained *female family* words (e.g., sister, mother). Similar to the findings in [15] and [20], we also observed that schizophrenia patients were more likely to use *feelings* words (especially feeling related to themselves), and less likely to use adverbs in their speech compared to control subjects.

This paper is organized as follows. In Section 2, we describe the experimental design and demographics of the participants. We then elaborate on the steps of our analysis in Section 3, and present the numerical results for speech recognition and linguistic analysis in Section 4. In Section 5, we investigate the most salient linguistic features, and offer our concluding remarks in Section 6.

## 2 Experiment Design

This experiment is in collaboration with the Institute of Mental Health Singapore (IMH). 71 individuals participated in this experiment. There are two groups of participants: 47 *Patients* who are diagnosed with schizophrenia, and 24 *Controls*, who do not have any pre-existing disorders. The participants are recruited by IMH based on the recommendation of clinicians, and the participants are matched for age, gender, ethnicity, and education. The participants are all above 19 years old, and although they are not native English speakers, Singapore being a multiracial and multicultural country, all of them can communicate in fluent English. The participants have provided written informed consent and receive monetary compensation for their participation in the study. The study protocol has been approved by the National Healthcare Group’s Domain-specific Review Board in Singapore. The demographic information of participants is displayed in Table 1.

In this experiment, each participant underwent an interview conducted by a professional psychometrician from IMH. The interview is a semi-structured one where the participants are asked certain fixed questions, but replies to those questions can lead to optional follow-up questions. These questions follow from the Negative Symptoms Assessment (NSA-16) rating instrument, which is 16-point scale specifically designed to reflect on the emotions and activities in the life of a patient suffering from negative symptoms of schizophrenia. Based on their replies, the psychometrician rates the behavior of the participants on the NSA-16 instrument on a scale of 1 to 6, where a rating of 1 denotes no recognizable symptoms and a rating of 6 denotes severe symptoms. There is no pre-determined time limit for the interview, nor role-playing during the interview. On average, the interviews last for around 25


		Patients (N = 47)	Healthy Controls (N = 24)
<b>Age</b>	Mean (years)	30.4	29.8
	Range (years)	20-49	19-47
<b>Gender</b>	Male	22	10
	Female	25	14
<b>Ethnicity</b>	Chinese	39	20
	Malay	5	3
	Indian	3	1
<b>Education</b>	University	6	4
	Diploma/JC/ITE	26	14
	High School	15	6

**Table 1** Demographics of participants.

minutes. We analyzed the entire length of the interview recordings. The total length of audio analyzed is about 34 hours.

### 3 System Overview

Fig. 1 depicts the different steps in our analysis.




**Fig. 1** Diagram of the analysis pipeline.

We employed portable equipment to record audio during the interview. Both the psychometrician and patient wore a lapel microphone, and the two audio streams were recorded by an H4n recorder. The two-channel audio recordings were saved in a single .wav file on a laptop. The psychometrician and the participant were seated about 2 meters apart, and this setting minimizes cross-talk from the psychometrician channel onto the participant channel.

#### 3.1 Speaker Diarization

The speech of participant and the psychometrician were recorded on separate channels, as mentioned earlier. Nevertheless, there is still some interference from the

psychometrician channel onto the participant channel. We apply automated speaker diarization to remove the psychometrician voice from the participant channel. As illustrate in Fig. 2, we extract binary sequences from both speech signals, indicating when the participant and the psychometrician respectively were speaking (0: not speaking; 1: speaking).



**Fig. 2** Illustration of speech preprocessing. We first reduce the interference between both audio channels. Channel 1 and 2 are the original signals, from which we derive binary sequences 1 and 2, indicating when the psychometrician and participant respectively are speaking. Next, we apply one-dimensional erosion and dilation to the binary vectors (shown at the bottom of the figure).

To improve the effectiveness of speech recognition, preprocessing is necessary. In order to obtain cohesive speech segments, we apply one-dimensional erosion and dilation to the binary sequence of the participant. Firstly, we dilate the binary sequence 2 by a one-second structuring element, which filled up small gaps (less than 1s) in one speech segment without filling up the adjacent two sentences. Next, we erode and dilate the binary sequence by a two-second structuring element. These steps reduce the noise and incorrect automated transcriptions. At last, we obtain the filtered speech signal, containing mostly speech from the participant, by multiplying the participant audio channel (channel 2) with the binary sequence associated with the participant (sequence 2).

### 3.2 *Speech Recognition*

The collected audio was then subjected to different speech-to-text software. We applied several speech-to-text APIs to the recorded audio files of the participants. Concretely, we tested seven speech recognition engines/APIs<sup>1</sup> were tested in our study:

<sup>1</sup> Zhang, A. (2017). Speech Recognition (Version 3.8) [Software]. Available from [https://github.com/Uberi/speech\\_recognition#readme](https://github.com/Uberi/speech_recognition#readme).

CMU Sphinx<sup>2</sup>, Google Speech Recognition<sup>3</sup>, Google Cloud Speech API<sup>4</sup>, Wit<sup>5</sup>, Microsoft Bing Voice Recognition<sup>6</sup>, Houndify API<sup>7</sup> and IBM Speech to Text<sup>8</sup>. In our tests, the Google Cloud Speech API performed the best (as shown in Section 4.1), hence we applied that API in our analysis (total transcribed words per recording:  $M = 709$ ).

### 3.3 Linguistic Analysis

Following speech-to-text conversion, we applied a dictionary-based method of Natural Language Processing. We counted the different types of words or phrases using Linguistic Inquiry and Word Count 2015 (LIWC). In the classification task, the linguistic extracted from the text were used as attributes in several supervised machine learning algorithms to classify the participants into *Patient* or *Control* groups. We used the Weka tool in JAVA [21] to perform classification with leave-one-out cross-validation. We tested the following classifiers: Support Vector Machine (SVM), Multilayer Perceptron (MLP), Logistic Regression (LR), Multinomial Nave Bayes and 1-Nearest Neighbor (1NN) [22]. Because each feature has specific semantic meaning, and not all features may be relevant to our task, feature selection becomes essential during the classification task. We applied three feature ranking methods: SVM-Recursive Feature Elimination (RFE), Relief Attribute Evaluator (Relief), and chi-Squared Attribute Evaluator [23]. For all combinations of the classifier, ranking method, and the number of top features, we selected the top features of the training data and classified the test data for each fold in cross-validation. We present the classification results for the three ranking methods in Section 4.2. In Section 5, we elaborate on the most salient linguistic features, since they give insight into characteristic behaviors of schizophrenia patients.

## 4 Results

### 4.1 Speech Recognition

We tested several speech recognition APIs to convert speech to text. Because no API has been specifically designed for Singaporean accented English, we measured the

---

<sup>2</sup> <https://cmusphinx.github.io/>

<sup>3</sup> <http://www.chromium.org/developers/how-tos/api-keys>

<sup>4</sup> <https://cloud.google.com/speech/>

<sup>5</sup> <https://wit.ai/>

<sup>6</sup> <https://azure.microsoft.com/en-us/services/cognitive-services/speech/>

<sup>7</sup> <https://www.houndify.com/>

<sup>8</sup> <https://console.bluemix.net/catalog/services/speech-to-text>

Speech API	Participant ID									mean	STD
	1	2	3	4	5	6	7	8	9		
<b>Google Cloud</b>	0.87	0.79	0.84	0.84	0.66	0.82	0.78	0.83	0.93	<b>0.82</b>	0.07
<b>Google Recognition</b>	0.85	0.83	0.84	0.71	0.62	0.77	0.68	0.81	0.83	0.77	0.08
<b>Bing</b>	0.68	0.67	0.62	0.7	0.42	0.72	0.32	0.68	0.69	0.61	0.14
<b>Houndify</b>	0.6	0.44	0.59	0.67	0.46	0.63	0.54	0.51	0.53	0.55	0.08
<b>IBM</b>	0.42	0.27	0.37	0.65	0.28	0.6	0.25	0.26	0.27	0.38	0.15
<b>Wit</b>	0.2	0.24	0.3	0.59	0.29	0.6	0.42	0.31	0.26	0.36	0.15
<b>Sphinx</b>	0.18	0.18	0.08	0.45	0.16	0.27	0.19	0.42	0.07	0.22	0.14

**Table 2** Transcription accuracies of different speech recognition APIs, tested on 9 random recordings in our dataset.

performance of seven speech recognition APIs on our dataset. We randomly chose 9 different audio files from our dataset, then extracted the first 5 minutes of each audio to test the accuracy of different speech recognition APIs. We determined the number of correct words in every transcription manually. We calculated the accuracies of translation of different APIs by dividing the number of correctly transcribed words by the number of total words (ground truth). The average transcription accuracy for each file and each API, and its standard deviation is listed in Table 2, where we transcribed on average 170 words in each file. These results indicate that Google Cloud speech API performed the best on our dataset, with an average accuracy of about 82% and standard deviation of around 7%. Therefore, we chose Google Cloud speech to convert all 71 recordings to text files.

## 4.2 Classification

In this paper, we tested several classifiers and feature ranking methods with the normalized LIWC features as attributes and class labels as targets. We present results for the three best-performing classifiers and their ranking method in Table 3. Both SVM and LR yielded the highest accuracy of 86%, which indicated the schizophrenia patients and controls in our dataset can be well separated by lexical features. We discuss the most salient linguistic features in Section 5.

## 5 Discussion

In Fig 3, we show how the classification accuracy varies with the number of linguistic features, for the best classifiers obtained by the three feature selection procedures. The SVM-RFE ranking method yields higher classification accuracy than the chi-square ranking method and leads to the same level of accuracy with fewer features compared to relief ranking method. Therefore, we selected the top 20 fea-

Classifier	Confusion Matrix		Precision	Recall	F-score	AUC	Accuracy	Baseline	Ranking method	
	P	C								
	<b>SVM</b>	<b>P</b>								45
	<b>C</b>	8	16	0.89	0.67	0.76	0.81			
<b>LR</b>	<b>P</b>	41	6	0.91	0.87	0.89	0.87	85.9%	66.2%	<b>SVM-RFE</b>
	<b>C</b>	4	20	0.77	0.83	0.80	0.87			
<b>1NN</b>	<b>P</b>	44	3	0.85	0.94	0.89	0.80	84.5%	66.2%	<b>chi-square</b>
	<b>C</b>	8	16	0.84	0.74	0.74	0.80			

**Table 3** Patient (P) vs. controls (C) classification with LIWC features.



**Fig. 3** Classification accuracy of three feature ranking methods as a function of the number of features. The results for the best performing classifiers for each feature selection approach are shown here.


Words patients used <i>less</i>			Words patients used <i>more</i>		
Category	Examples	p-value	Category	Examples	p-value
<b>Informal</b>	netspeak, swear words	0.0029	<b>Female family</b>	lady, sister, mom	0.0024
<b>Netspeak</b>	btw, lol, thx	0.0046	<b>Feel</b>	feels, touch, anger	0.0602
<b>Assent</b>	agree, ok, yes	0.0056	<b>Family</b>	mom, father, wife	0.0935
<b>Adverbs</b>	very, really, quickly	0.0090			

**Table 4** Word categories showing with different average counts for patients and healthy controls.

tures by means of the SVM-RFE. Next, we apply the Kruskal-Wallis Test to these 20 linguistic features, and computed the corresponding p-values. The resulting most salient categories and their p-values are shown in Table 4.

We separated the categories into two classes: the type of words schizophrenic patients used *less* and *more* respectively compared to control subjects. The box plots of the four most salient categories are shown in Fig. 4. We found that patients were less likely to use *adverb* words and *informal* words, which include the *assent* category and *netspeak* category. On the other hand, we found that *female family* words

and *feel* words appeared more frequently in the speech produced by schizophrenic patients compared to controls.



**Fig. 4** Box plots of *female family*, *informal*, *assent*, and *feel* categories for patients and healthy controls.

We computed the correlation coefficient for *I* and *feeling*, and for *family* and *feeling*; we also conducted the Pearson correlation test. The results of this analysis are summarized in Tabel 5. We can see from this table that schizophrenia patients tend to use both the word *I* and words related to feelings simultaneously more often compared to control subjects. This effect is also present, yet less pronounced, for the simultaneous usage of *family* and *feeling* words. These speech pattern results appear to be in line with the research on social cognitive impairments of schizophrenic patients [26]. Studies have found that due to impairment in social cognition [24, 25], schizophrenic patients tend to focus on themselves, their feelings and thoughts in a conversation, reflecting an inability to perceive, interpret and generate responses to the intentions, dispositions, and behaviors of others [20]. In [15], it was observed that patients were more likely to focus on their own feelings when writing autobiographic narratives as compared to healthy controls.

We also found in our study that the *family* words were more frequently mentioned by schizophrenic patients. It is plausible that for patients, their family members are often the ones they interact with the most within their social circles, outside of therapeutic environments. Thus, the frequent mention of *family* is plausible when referencing themselves in the conversation.

Studies that investigate differences in LIWC categories in schizophrenic patients and healthy controls often make use of the LIWC 2007 version. *Female* and *informal* are a new category in the 2015 version that we applied here. Therefore, it would be hard to make direct comparisons of the results of this study with earlier studies. In our study, we observed that female patients used far more *female family* vocabulary (mean=0.39 per minute) than healthy female controls (mean=0.1 per minute)

	<b>Feeling &amp; I</b>		<b>Feeling &amp; Family</b>	
	<b>Correlation</b>	<b>p-value</b>	<b>Correlation</b>	<b>p-value</b>
<b>Patients</b>	0.648	8.66E-07	0.287	0.05
<b>Healthy</b>	0.5	0.0129	0.026	0.905

**Table 5** Correlation coefficients and p-values of the Pearson correlation test for *feeling vs. I* and *feeling vs. family*.

<b>Female family words (per min)</b>				<b>Male family words (per min)</b>			
		<b>Mean</b>	<b>STD</b>			<b>Mean</b>	<b>STD</b>
<b>Patient</b>	<b>Female</b>	<b>0.39</b>	0.37	<b>Patient</b>	<b>Female</b>	0.22	0.14
	<b>Male</b>	0.17	0.11		<b>Male</b>	0.15	0.08
<b>Healthy</b>	<b>Female</b>	<b>0.10</b>	0.13	<b>Healthy</b>	<b>Female</b>	0.22	0.32
	<b>Male</b>	0.16	0.16		<b>Male</b>	0.21	0.10

**Table 6** Word frequency of *female family* and *male family* category on patient and healthy people.

	<b>Female family words</b>	<b>Male family words</b>
<b>Female patients</b>	<b>0.00055</b>	0.250
<b>Male patients</b>	0.515	0.046

**Table 7** Patients v/s Controls p-values of *female family* words and *male family* words.

as shown in Table 6. This trend did not show up between male patients and healthy male controls. In Table 7, we further confirm that only female patients used more *female family* words ( $p=0.00055$ ).

The existing literature suggests that female schizophrenic patients often report higher levels of social support, social functioning, and social withdrawal than male patients [27, 28]. Traditionally, caretaking roles often fall to female members of the family [29], and female caretakers of patients perceive caretaking of their schizophrenic family members more rewarding as well [30]. Consequently, female patients may have greater access and higher proximity to a social support network of female caretakers (aunt, mother, sister), which could account for the higher occurrence of *female family* words in their speech.

## 6 Conclusion

In this paper, we analyzed linguistic features extracted from interviews of schizophrenic patients and healthy control subjects. More precisely, we applied LIWC2015 to provide a dictionary-based words counting method for extracting linguistic features from the audio recordings. We applied several classification and feature ranking

methods to distinguish patient and healthy subjects. We obtained an accuracy of 86% for distinguishing schizophrenia patients from healthy control subjects on our dataset of 71 participants. We observed that patients used *informal* and *adverb* words less frequently, but instead used more words correlated to *female family* and *feel*. However, the results are limited by the accuracy of speech recognition, since the accuracy of converting Singapore English to text is significantly lower than for native English (US and UK). Moreover, the sample size (47 patients and 24 control subjects) is relatively small. More research is warranted to further explore and demonstrate the results of this study. In future work, we will also explore how combining non-verbal and verbal features may provide a more comprehensive characterization of schizophrenia patients.

**Acknowledgements** This study was funded by the Singapore Ministry of Health National Medical Research Council Center Grant awarded to the Institute of Mental Health Singapore (NMRC/CG/004/2013) and by NITHM grant M4081187.E30. This research is also supported in part by the Being Together Centre, a collaboration between Nanyang Technological University (NTU) Singapore and University of North Carolina (UNC) at Chapel Hill. The Being Together Centre is supported by the National Research Foundation, Prime Minister's Office, Singapore under its International Research Centres in Singapore Funding Initiative. Besides, this project is funded in part by the RRIS Rehabilitation Research Grant RRG2/16009. The authors also acknowledge support from the Interdisciplinary Graduate School at NTU.

## References

1. Demily C, Franck N. Cognitive remediation: a promising tool for the treatment of schizophrenia. *Expert Review of Neurotherapeutics*. 2008 Jul 1;8(7):1029-36.
2. Covington MA, He C, Brown C, Nai L, McClain JT, Fjordbak BS, Semple J, Brown J. Schizophrenia and the structure of language: the linguist's view. *Schizophrenia research*. 2005 Sep 1;77(1):85-98.
3. Andreasen NC. Thought, language, and communication disorders: I. Clinical assessment, definition of terms, and evaluation of their reliability. *Archives of general Psychiatry*. 1979 Nov 1;36(12):1315-21.
4. Slatcher RB, Trentacosta CJ. Influences of parent and child negative emotionality on young children's everyday behaviors. *Emotion*. 2012 Oct;12(5):932.
5. Rude S, Gortner EM, Pennebaker J. Language use of depressed and depression-vulnerable college students. *Cognition & Emotion*. 2004 Dec 1;18(8):1121-33.
6. Losh M, Gordon PC. Quantifying narrative ability in autism spectrum disorder: A computational linguistic analysis of narrative coherence. *Journal of autism and developmental disorders*. 2014 Dec 1;44(12):3016-25.
7. Jackson RG, Patel R, Jayatilleke N, Kolliakou A, Ball M, Gorrell G, Roberts A, Dobson RJ, Stewart R. Natural language processing to extract symptoms of severe mental illness from clinical text: the Clinical Record Interactive Search Comprehensive Data Extraction (CRIS-CODE) project. *BMJ open*. 2017 Jan 1;7(1):e012012.
8. Thinking big in mental health. *Nature Medicine*. 2018 Jan 09; 24(1):1. Nature Publishing Group.
9. Tausczik YR, Pennebaker JW. The psychological meaning of words: LIWC and computerized text analysis methods. *Journal of language and social psychology*. 2010 Mar;29(1):24-54.
10. Pennebaker JW, Boyd RL, Jordan K, Blackburn K. The development and psychometric properties of LIWC2015. 2015 Sep 15.

11. St-Hilaire A, Cohen AS, Docherty NM. Emotion word use in the conversational speech of schizophrenia patients. *Cognitive neuropsychiatry*. 2008 Jul 1;13(4):343-56.
12. Jungthaenel DU, Smyth JM, Santner L. Linguistic dimensions of psychopathology: A quantitative analysis. *Journal of Social and Clinical Psychology*. 2008 Jan;27(1):36-55.
13. Deutsch-Link S. Language In Schizophrenia: What We Can Learn From Quantitative Text Analysis.
14. Minor KS, Bonfils KA, Luther L, Firmin RL, Kukla M, MacLain VR, Buck B, Lysaker PH, Salyers MP. Lexical analysis in schizophrenia: how emotion and social word use informs our understanding of clinical presentation. *Journal of psychiatric research*. 2015 May 31;64:74-8.
15. Hong K, Kohler CG, March ME, Parker AA, Nenkova A. Lexical differences in autobiographical narratives from schizophrenic patients and healthy controls. In *Proceedings of the 2012 Joint Conference on Empirical Methods in Natural Language Processing and Computational Natural Language Learning 2012 Jul 12* (pp. 37-47). Association for Computational Linguistics.
16. Tahir Y, Chakraborty D, Dauwels J, Thalmann N, Thalmann D, Lee J. Non-verbal speech analysis of interviews with schizophrenic patients. In *Acoustics, Speech and Signal Processing (ICASSP), 2016 IEEE International Conference on 2016 Mar 20* (pp. 5810-5814). IEEE.
17. Chakraborty D, Tahir Y, Yang Z, Maszczyk T, Dauwels J, Thalmann D, Thalmann NM, Tan BL, Lee J. Assessment and prediction of negative symptoms of schizophrenia from RGB+ D movement signals. In *Multimedia Signal Processing (MMSP), 2017 IEEE 19th International Workshop on 2017 Oct 16* (pp. 1-6). IEEE.
18. Lange P, Suendermann-Oeft D. Tuning Sphinx to Outperform Googles Speech Recognition API. In *Proc. of the ESSV 2014, Conference on Electronic Speech Signal Processing 2014* (pp. 1-10).
19. Hinton G, Deng L, Yu D, Dahl GE, Mohamed AR, Jaitly N, Senior A, Vanhoucke V, Nguyen P, Sainath TN, Kingsbury B. Deep neural networks for acoustic modeling in speech recognition: The shared views of four research groups. *IEEE Signal Processing Magazine*. 2012 Nov;29(6):82-97.
20. Green MF, Penn DL, Bentall R, Carpenter WT, Gaebel W, Gur RC, Kring AM, Park S, Silverstein SM, Heinsen R. Social cognition in schizophrenia: an NIMH workshop on definitions, assessment, and research opportunities. *Schizophrenia bulletin*. 2008 Jan 8;34(6):1211-20.
21. Witten IH, Frank E, Trigg LE, Hall MA, Holmes G, Cunningham SJ. Weka: Practical machine learning tools and techniques with Java implementations.
22. Hall M, Frank E, Holmes G, Pfahringer B, Reutemann P, Witten IH. The WEKA data mining software: an update. *ACM SIGKDD explorations newsletter*. 2009 Nov 16;11(1):10-8.
23. Frank E, Hall M, Holmes G, Kirkby R, Pfahringer B, Witten IH, Trigg L. Weka-a machine learning workbench for data mining. In *Data mining and knowledge discovery handbook 2009* (pp. 1269-1277). Springer, Boston, MA.
24. Penn DL, Corrigan PW, Bentall RP, Racenstein J, Newman L. Social cognition in schizophrenia. *Psychological bulletin*. 1997 Jan;121(1):114.
25. Savla GN, Vella L, Armstrong CC, Penn DL, Twamley EW. Deficits in domains of social cognition in schizophrenia: a meta-analysis of the empirical evidence. *Schizophrenia bulletin*. 2012 Sep 4;39(5):979-92.
26. Howes C, Purver M, McCabe R. Using conversation topics for predicting therapy outcomes in schizophrenia. *Biomedical informatics insights*. 2013;6(Suppl 1):39.
27. Hfner H. Gender differences in schizophrenia. *Psychoneuroendocrinology*. 2003 Apr 30;28:17-54.
28. Ochoa S, Usall J, Cobo J, Labad X, Kulkarni J. Gender differences in schizophrenia and first-episode psychosis: a comprehensive literature review. *Schizophrenia research and treatment*. 2012 Apr 8;2012.
29. Diekman AB, Goodfriend W. Rolling with the changes: A role congruity perspective on gender norms. *Psychology of Women Quarterly*. 2006 Dec 1;30(4):369-83.
30. Chen FP, Greenberg JS. A positive aspect of caregiving: The influence of social support on caregiving gains for family members of relatives with schizophrenia. *Community mental health journal*. 2004 Oct 1;40(5):423-35.